

Collective quantities of a fermionic particle beam in a circular accelerator near zero temperature

Andreas Kabel*

*Stanford Linear Accelerator Center,
2575 Sand Hill Road, Menlo Park, CA 94025
E-Mail: andreas.kabel@slac.stanford.edu*

In the usual parameter regime of accelerator physics, particle ensembles can be treated as classical. If we approach a regime where $\epsilon_x \epsilon_y \epsilon_s \approx N_{\text{particles}} \lambda_{\text{Compton}}^3$, however, the granular structure of quantum-mechanical phase space becomes a concern. In particular, we have to consider the Pauli exclusion principle, which will limit the minimum achievable emittance for a beam of fermions. We calculate these lowest emittances for the cases of bunched and coasting beams at zero temperature and their first-order change rate at finite temperature. The self-field of the particle beam will, analogous to a space-charge tune depression, lead to a decrease of the quantum-mechanical state density. We calculate the tunes and the increased emittance for this case.

I. INTRODUCTION

When a relativistic particle travels in a focusing dominated beamline, it can be viewed, in its rest frame, as a particle in a harmonic oscillator potential. Emission of dipole radiation will lead to transverse energy loss; quantum-mechanically, the particle will drop down one energy level by emission of a photon. This mechanism will, in the absence of competing heating mechanisms, ultimately lead to a cooling down of the particle's transverse action to its quantum-mechanical limit of $J_{\text{min}} = \frac{\hbar}{2}$. This result has been established in [1], however, only single-particle dynamics is considered there.

In the multi-particle case, some care has to be taken if the particles in question are fermions, as Pauli's exclusion principle has to be considered when constructing the ground state; it has been claimed elsewhere[2] that this will limit the admissible minimum emittance to values not too far out of the reach of today's technologies.

In this paper, the emittances for a relativistic fermion bunch and a coasting beam are calculated for temperatures close to zero. We will show that the quantum-mechanical limit for the transverse emittances for typical configurations are vastly smaller than values reached today.

II. CALCULATING COLLECTIVE QUANTITIES: THE WIGNER TRANSFORMATION

In this paper, we want to calculate the emittance of a multi-particle system. The classical emittance is defined in terms of classical expectation values, i. e. particle averages, as $\epsilon^2 = \langle p^2 \rangle_{\text{Cl}} \langle q^2 \rangle_{\text{Cl}} - \langle pq \rangle_{\text{Cl}}^2$. A quantum version of this and similar quantities can be obtained by interpreting the Wigner transform of the state of the quantum system in question as a classical probability distribution.

Consider an N -particle system in a pure state $|\Psi\rangle = P|n_1\rangle \cdots |n_n\rangle$, where P is the (anti)symmetrizing operator, the particles in question being (fermions) bosons. Now, we can create a pseudo-classical probability distribution $\rho(q_1, p_1; \dots; q_N, p_N)$ by using the Wigner transform of the product wavefunction. It is easy to see that all expectation values of this phase-space distribution are the same as the ones of the Wigner transform of a one-particle statistical matrix with equal weights in all states. We can thus conclude that the appropriate quantum version of the classical expectation values above is obtained by replacing the classical average $\langle f(p, q) \rangle$ by $\langle \langle f(p_{op}, q_{op}) \rangle_{\text{Qm}} \rangle_{|n_i\rangle}$, i. e., the quantum-mechanical expectation value for the corresponding operator function, averaged over the occupied states.

III. DYNAMICS

Consider an ultra-relativistic particle beam in a circular accelerator. Neglecting higher-order effects, the Hamiltonian can be written as a quadratic form in the usual phase-space coordinates $x, x', y, y', \sigma, \delta$.

*Work supported by Department of Energy contract DE-AC03-76SF00515.

However, this Hamiltonian is not appropriate for quantization, as energy and time have switched roles. Thus, we use the Hamiltonian of the system in the beam's frame of reference, which can be obtained by a series of canonical transformation from the lab frame[3]:

$$H = \frac{p_x^2}{2} + \frac{p_y^2}{2} + \frac{p_z^2}{2} - \frac{\gamma^2 \beta x p_z}{R} + \beta^2 \gamma^2 \left(\kappa_x - \frac{\gamma^2}{R^2} \right) \frac{x^2}{2} + \beta^2 \gamma^2 \kappa_y \frac{y^2}{2} + \Phi_{RF}(z) \quad , \quad (1)$$

where γ is the relativistic factor, $\kappa_{x,y}$ the (possibly local) focusing strengths (in the case of magnetic quadrupoles, one has $\kappa_x = -\kappa_y$) and $\Phi(z)$ the external electric potential. The directions x, y, z are radial, transversal, and tangential, respectively. Note that we use units with $\hbar = c = k_B = m_0$ throughout, so all quantities are expressed in powers of the Compton length of the particles in question.

The longitudinal part of the Hamiltonian depends on the physical setup. The particles might either be confined by the nearly harmonic potential of the RF bucket, or we have case of a coasting beam, where the only constraints imposed on the longitudinal motion are the ones due to the periodicity of the problem. In the sequel, we will consider both cases.

IV. ANISOTROPIC OSCILLATOR

Let us assume that the longitudinal motion is determined by the presence of an RF bucket. We can approximate the potential Φ by expanding it to 2nd order in z . For reasons of simplicity, we only take into account the $O(z^2)$ term, i.e., we assume that the particle is on the orbit and is not losing energy.

For a bunched beam with dimensions σ_z, σ_δ , we have, in the orbit frame, $\omega_l \gamma \sigma_l = \sigma_\delta$.

The longitudinal and radial parts of the hamiltonian (1) have the form

$$H = \sum_{i=1}^2 \frac{p_i^2}{2} + \frac{\omega_i^2 q_i^2}{2} - \mu q_1 p_2 \quad . \quad (2)$$

Its eigenfrequencies are determined by the equation

$$\lambda^4 + \lambda^2(\omega_1^2 + \omega_2^2) + \omega_2^2(\mu^2 - \omega_1^2) = 0 \quad , \quad (3)$$

which leads to stable motion for

$$(\omega_1^2 + \omega_2^2) > 4\omega_2^2(\mu^2 - \omega_1^2) > 0 \quad .$$

The first condition will always be fulfilled for realistic machines. The second one corresponds to the machine being below or above transition: if the second factor changes sign, the eigenfrequency can be made real again by flipping the sign of $\omega_z^2 \propto \Phi''_{RF}$. However, the absolute sign of both the kinetic and the potential term will change, leading to the (for purposes of constructing the quantum-mechanical ground state) pathological case of a hamiltonian not limited from below. In the sequel, we will assume the machine is below transition.

Thus, expanding (1) to first non-trivial order in the canonical coordinates and applying the canonical transformation removing the mixed term in (2), we obtain the Hamiltonian of a 3-dimensional harmonic oscillator with corrected frequencies given by (3); the ground state is characterized by the occupation numbers $n_n^d \in \{0, 1\}$ where $\sum n_d^i = N$ of the oscillator levels. For sake of generality, we consider the case of d dimensions. The ground state for a given particle number can be constructed by successively filling states with the lowest energy (we disregard spin here, which can be easily reintroduced by replacing $N \rightarrow 2N$ in the final formulae).

In $\frac{E}{\epsilon_F}$ -space, the Fermi sea is just a unit d -simplex, in n_i -space, a d -simplex with axes of length $\frac{\omega_1}{\epsilon_F}, \dots, \frac{\omega_d}{\epsilon_F}$. Thus, the particle number for a ground state filled up to the Fermi energy ϵ_F , where we have disregarded the zero-mode energy $\frac{1}{2} \sum_i \omega_i$ of the oscillator,

$$N = \frac{\epsilon_F^d}{\Omega d!} \quad ,$$

the volume of an unit d -simplex being $\frac{1}{d!}$ and $\Omega = \omega_1 \cdots \omega_d$.

The energy in the i th degree of freedom in that case is given by a sum over the d -simplex. Replacing all sums by integrals, we have

$$E_i = \frac{\epsilon_F^d}{\Omega} \int_0^1 \int_0^{1-q_1} \cdots \int_0^{1-\sum_{k=1}^{d-1} q_k} \epsilon_F q_i dq_1 \cdots dq_d = \frac{\epsilon_F^{d+1}}{\Omega(d+1)!} = \frac{N \epsilon_F}{d+1} \quad .$$

We calculate the emittance by using its statistical definition $\varepsilon^2 = \langle p^2 \rangle q^2 - \langle pq \rangle^2$ and replacing the classical averages by the ones discussed in II.

For a harmonic oscillator, we have $\langle p^2 \rangle_{\text{Qm}} \langle q^2 \rangle_{\text{Qm}} = n$ and $\langle pq \rangle_{\text{Qm}} = 0$, so:

$$\varepsilon_i = \langle n_i \rangle_{|n_i\rangle} = \frac{E_i}{N\omega_i} = \frac{\epsilon_F}{\omega_i(d+1)} = \frac{\Omega^{\frac{1}{d}}}{\omega_i(d+1)} \sqrt[d]{Nd!} \quad . \quad (4)$$

and the total phase-space volume

$$\begin{aligned} \varepsilon^{(d)} &= \prod_{i=1}^d \varepsilon_i = \frac{Nd!}{(d+1)^d} \\ \varepsilon^{(3)} &= \frac{3}{32}N \end{aligned}$$

Thus, the projected emittances scale as $N^{\frac{1}{d}}$, as one would naïvely assume. Furthermore, due to the occurrence of the geometric mean of the frequencies in (4), the projected emittance in one dimension can be lowered by shallowing the potential in the other dimensions.

Note that a similar approach has been chosen elsewhere; [[2]] gives an estimate for ε_{min} from a similar reasoning, but ends up (due to a miscounting of the states) with a scaling different from our result.

V. MIXED CASE: LONGITUDINALLY FREE PARTICLES

So far, we have assumed an anisotropic oscillator. But given the case of a particle moving freely longitudinally, the energy content of that degree of freedom will be given by the square of the (angular) momentum. (We might consider the boundary conditions imposed by a periodic box instead of a circular arrangement.) We treat the general case, i. e., a hamiltonian

$$H = \sum_{\tilde{i}=1}^{\tilde{d}} \tilde{\omega}_{\tilde{i}}^2 \tilde{n}_{\tilde{i}}^2 + \sum_{i=1}^d \omega_i n_i \quad .$$

The emittance in each of $\tilde{d}$ new degrees of freedom is given by $\langle p^2 \rangle_{\text{Qm}} = \frac{n^2 \pi^2}{D^2} = 2E_n$ and $\langle q^2 \rangle_{\text{Qm}} = \frac{D^2}{12}$, where D is the length of the enclosing potential, and $\langle pq \rangle_{\text{Qm}} = 0$, so

$$\tilde{\varepsilon}_{\tilde{i}} = D \sqrt{\frac{\langle \tilde{E}_{\tilde{i}} \rangle_{|n\rangle}}{6}} = \sqrt{\langle \tilde{E}_{\tilde{i}} \rangle_{|n\rangle}} \frac{\pi}{\sqrt{12} \tilde{\omega}_{\tilde{i}}} \quad .$$

Rescaling the integration range to a unit sphere and unit simplex, we get

$$N = \frac{\epsilon_F^{\frac{\tilde{d}}{2}+d}}{\tilde{\Omega}\Omega} \int_{\tilde{d}\text{-sphere}} \int_{d\text{-simplex}}^{1-\tilde{q}^2} dq d\tilde{q} = \frac{\pi^{\frac{\tilde{d}}{2}} \epsilon_F^{\frac{\tilde{d}}{2}+d}}{\tilde{\Omega}\Omega \Gamma\left(\frac{2d+\tilde{d}+2}{2}\right)} \quad , \quad (5)$$

We can readily write down the averaged values of the energy in the different degrees of freedom:

$$\frac{\langle \tilde{E}_{\tilde{i}} \rangle}{\epsilon_F} = \langle \tilde{q}^2 \rangle = \frac{\int_0^1 \tilde{q}^2 (1-\tilde{q}^2)^{\frac{\tilde{d}-1}{2}+d} d\tilde{q}}{\int_0^1 (1-\tilde{q}^2)^{\frac{\tilde{d}-1}{2}+d} d\tilde{q}} = \frac{1}{2d+\tilde{d}+2}$$

and

$$\frac{\langle E_i \rangle}{\epsilon_F} = \langle q \rangle = \frac{\int_0^1 q (1-q)^{\frac{\tilde{d}}{2}+d-1} dq}{\int_0^1 (1-q)^{\frac{\tilde{d}}{2}+d-1} dq} = \frac{2}{2d+\tilde{d}+2} \quad (6)$$

so

$$\varepsilon_i = \frac{\epsilon_F}{\omega_i} \cdot \frac{2}{\tilde{d}+2d+2} \propto N^{\frac{2}{\tilde{d}+2d}} \quad (7)$$

and

$$\tilde{\varepsilon}_i = \sqrt{\frac{\epsilon_F}{\tilde{\omega}_i}} \cdot \frac{\pi}{\sqrt{12}(\tilde{d} + 2d + 2)} \propto N^{\frac{1}{d+2d}}$$

and the product emittance is

$$\varepsilon^{(d,\tilde{d})} = \frac{\pi^{\tilde{d}} 2^d \epsilon_F^{d+\frac{d}{2}}}{12^{\frac{d}{2}} (\tilde{d} + 2d + 2)^{d+\frac{\tilde{d}}{2}} \Omega \tilde{\Omega}} = \left(\frac{\pi}{24}\right)^{\frac{\tilde{d}}{2}} \cdot \frac{\Gamma\left(\frac{\tilde{d}}{2} + d + 1\right)}{\left(\frac{\tilde{d}}{2} + d + 1\right)^{\frac{\tilde{d}}{2}+d}} N \quad .$$

Of course, by putting $\tilde{d} = 0$, we regain the formulae for the bunched-beam case.

Putting in a real coasting-beam ring, we can express the orbit-frame frequencies by the tune[4]: $\omega_x \approx \omega_y = \frac{\beta\nu_y}{\gamma L}$, where L is the length of the ring. The longitudinal momentum is quantized in units of $\frac{\pi}{\gamma L}$, so $\omega_l = \frac{\pi}{\sqrt{2}\gamma L}$ and

$$\varepsilon_x = \langle q \rangle \frac{\epsilon_F}{\omega_x} = \frac{1}{7} \sqrt[5]{\frac{225\pi N^2}{256\gamma L\nu}} \quad .$$

This means the transverse degrees of freedom begin to exhibit a non-minimal emittance for $N > N_0$, where

$$N_0 \approx 78.0 \sqrt{\frac{\nu\gamma L}{\lambda_{\text{Compton}}}}$$

in usual units, and the transverse emittances will grow $\propto N^{\frac{2}{5}}$ for higher particle numbers.

For parameters one could consider for a real focusing-dominated, below-transition ring ($\gamma = 10, \nu = 100, L = 2\pi\text{m}$), we get $N_0 \approx 1.0 \cdot 10^{10}$ particles; i.e., the beam will have, for realistic particle numbers, an emittance close to its minimum value. (λ_{Compton} in customary units.)

VI. SELF-CONSISTENT TUNE SHIFT FOR THE FERMI CONDENSATE

In our construction, we tacitly assume that the particle-particle interaction does not modify the particle content of the ground state. This corresponds precisely to the notion of a Fermi liquid (in our case, a highly anisotropic one), in which the free particle spectrum smoothly deforms into the quasi-particle spectrum of equal particle content when the interaction is switched on adiabatically. This naïve assumption of the existence of a Fermi surface may break down if we take into account particle-particle interactions.

For the case of the system being below transition, we can make the following semi-quantitative argument for the existence of a Fermi liquid: In a “mean-field calculation”, we estimate the effective transverse focusing strength ω_{eff} to be the sum of the external focusing and a space-charge tune depression due to a circular beam of radius $\sqrt{\langle x^2 \rangle}$:

$$\omega_{\text{eff}}^2 = \omega_{\text{ext}}^2 - \omega_{\text{SC}}^2 = \omega_{\text{ext}}^2 - \frac{Ne^2}{2\gamma L \langle x \rangle^2}.$$

In the following, we consider a coasting beam. We have $\langle x^2 \rangle \omega_{\text{eff}}^2 = \langle E_x \rangle$, where $\langle E_x \rangle$ is given by (6). We use (5) with $\omega = \omega_{\text{eff}}$ to eliminate ϵ_F and obtain a consistency condition for ω_{eff} :

$$\left(\frac{\omega_{\text{eff}}}{\omega_{\text{ext}}}\right)^2 \left[1 + \left(\frac{\omega_{\text{int}}}{\omega_{\text{ext}}}\right)^{4/5} \left(\frac{\omega_{\text{eff}}}{\omega_{\text{ext}}}\right)^{-4/5}\right] = 1 \quad , \quad (8)$$

where we have introduced

$$\omega_{\text{int}} = \omega_{\text{ext}} \frac{\sqrt[4]{2^3 7^5 r_{\text{cl}}^3 (\gamma L) N^3}}{\nu \sqrt{15\pi^3 \lambda_{\text{Compton}}^3}} \quad . \quad (9)$$

(8) has a solution $\omega_{\text{eff}} < \omega_{\text{ext}}$ for all N . This corrected frequency has to be substituted in the equations in the previous section, leading to corrected values for the emittances.




FIG. 1: Effective focusing strength of a coasting beam in its ground state; $\omega_{int} = e^3 N / (\gamma L)$

For typical setups, $\omega_{int} \gg \omega_{ext}$

(using the ring parameters from above and $N = 10^{10}$ particles, we end up with $\omega_{int}/\omega_{ext} \approx 2.14 \cdot 10^6$ and $\varepsilon_{i,corrected} \approx 18.5\varepsilon_i$), so $\frac{\omega_{eff}}{\omega_{ext}} \approx \left(\frac{\omega_{int}}{\omega_{ext}}\right)^{-\frac{2}{3}}$. The scaling of the transverse emittances is $\propto \omega_{eff}^{\frac{1}{5}}$; with (9), $\omega_{eff} \propto N^{-\frac{1}{2}}$, so we end up with a corrected scaling for the case of large particle numbers: $\varepsilon_i \propto N^{\frac{1}{2}}$ instead of $\varepsilon_i \propto N^{\frac{2}{5}}$ from (7).

This modified scaling law means that the transverse phase space volume scales as $\varepsilon_x \varepsilon_y = O(N)$: adding a new particle to the coasting beam's ground state will lower the transverse frequencies, thus increasing the state density, so a new site becomes available in transverse phase space, while the longitudinal phase space retains its original volume. In particular, this means that kinetic and potential energy always will balance out, $\epsilon_F \approx \epsilon_{kin} \approx \epsilon_{pot}$, keeping the system at the boundary of a “weakly non-ideal” and a “strongly non-ideal” gas. Thus, the simple criterion $\epsilon_{kin} \ll \epsilon_{pot}$ for quantum Wigner crystallization is not applicable for any external focusing strength. This property is peculiar to the external oscillator potential and is not present for other confining potentials.

VII. FINITE TEMPERATURE

The above considerations were for the case of zero temperature. To generalize to finite temperatures, we follow the usual prescription and introduce a chemical potential. The quantity we want to calculate is the logarithm of the partition function of the grand-canonical ensemble:

$$\log Z = \log \sum_{n_i \in \{0,1\}} e^{-\beta \sum_i n_i (\sum_{k=1}^d \omega_k (i_k + \frac{1}{2}) - \mu)} = \sum_{i_k=0}^{\infty} \log \left(1 + e^{-\beta (\sum \omega_k (i_k + \frac{1}{2}) - \mu)} \right) \quad .$$

Again, we transform the sum into an integral. With (5), we integrate over energy

$$\log Z = \frac{1}{\Omega \tilde{\Omega} \Gamma \left(d + \frac{d}{2} \right)} \int_0^{\infty} \log \left(1 + e^{-\beta (E - \mu)} \right) E^{d + \frac{d}{2} - 1} dE \quad .$$

For small temperatures, integrals of this style can be done by integrating by parts and then be approximated using a Sommerfeld expansion. We find

$$\log Z = -\frac{\beta\pi^{\frac{d}{2}}}{\Gamma\left(d+\frac{\tilde{d}}{2}+1\right)}\left(\frac{\mu^{d+1}}{d+\frac{\tilde{d}}{2}+1}+\left(d+\frac{\tilde{d}}{2}\right)\frac{\mu^{d-1}\pi^2}{6\beta^2}+\dots\right) \quad .$$

As

$$\left. \begin{aligned} \varepsilon_i &= -\frac{1}{\beta N} \frac{\partial}{\partial \omega_i} \log Z \\ \tilde{\varepsilon}_i &= \sqrt{-\frac{\pi^2}{24\beta N \omega_i} \frac{\partial}{\partial \tilde{\omega}_i} \log Z} \end{aligned} \right\} \quad ,$$

we can write the temperature-dependent contributions to the emittances:

$$\left. \begin{aligned} \frac{\Delta \varepsilon_i}{\varepsilon_i} &= \frac{T^2 \pi^2}{6\epsilon_F^2} \left(d + \frac{\tilde{d}}{2}\right) \left(d + \frac{\tilde{d}}{2} + 1\right) \\ \frac{\Delta \tilde{\varepsilon}_i}{\tilde{\varepsilon}_i} &= \frac{T^2 \pi^2}{12\epsilon_F^2} \left(d + \frac{\tilde{d}}{2}\right) \left(d + \frac{\tilde{d}}{2} + 1\right) \end{aligned} \right\} \quad (10)$$

where we have used the zero-temperature ϵ_F as chemical potential, which is correct in this order of β . Thus, the emittances grow quadratically in the temperature, with the scale being set by ϵ_F . For the parameters from above and $N = 10^{10}$ particles, one finds, from (10), a transverse emittance-doubling temperature of $T = 0.03K$.

VIII. ACKNOWLEDGMENTS

I wish to thank R. Ruth and M. Venturini for useful discussions.

-
- [1] Z. Huang, Ph.D. thesis, Stanford University (1999).
 - [2] A. A. Mikhailichenko, in *Proceedings of the 1999 Particle Accelerator Conference, New York* (1999).
 - [3] J. Wei, X.-P. Li, and A. M. Sessler, BNL Preprint, BNL-52381, UC-414 (1993).
 - [4] M. Venturini and R. Ruth, in *Proceedings of the Conference on Quantum Aspects of Beam Physics, Capri, Italy, 15-20 October 2000* (2000).